

Accepted for publication in the Letters of the Astrophysical Journal

The Bright SHARC Survey: The X-ray Cluster Luminosity Function ¹

R. C. Nichol & A. K. Romer

Dept. of Physics, Carnegie Mellon University, 5000 Forbes Avenue, Pittsburgh, PA-15213, USA.
(nichol@andrew.cmu.edu & romer@astro.phys.cmu.edu)

B. P. Holden

Dept. of Astronomy & Astrophysics, University of Chicago, 5640 S. Ellis Avenue, Chicago, IL-60637, USA.
(holden@oddjob.uchicago.edu)

M. P. Ulmer, R. A. Pildis & C. Adami

Dept. of Physics & Astronomy, Northwestern University, Dearborn Observatory, 2131 N. Sheridan Road,
Evanston, IL-60208, USA. (m-ulmer2@nwu.edu, pildis@enteract.com, adami@lilith.astro.nwu.edu)

A. J. Merrelli

Dept. of Physics, Carnegie Mellon University, 5000 Forbes Avenue, Pittsburgh, PA-15213, USA.
(merrelli@andrew.cmu.edu)

D. J. Burke & C. A. Collins

Astrophysics Research Institute, School of Engineering, Liverpool John Moores University, Twelve Quays
House, Egerton Wharf, Birkenhead, L41 1LD, UK. (djb@astro.livjm.ac.uk, cac@astro.livjm.ac.uk)

ABSTRACT

We present here initial results on the X-ray Cluster Luminosity Function (XCLF) from the Bright SHARC sample of distant X-ray clusters of galaxies. This sample is 97% complete in its optical identifications and contains 12 X-ray luminous clusters in the redshift range $0.3 \leq z \leq 0.83$ (median $z = 0.42$) and $1.1 \leq L_x \leq 8.5 \times 10^{44} \text{ erg s}^{-1}$ [0.5–2.0 keV]. We present a preliminary selection function for the Bright SHARC based on Monte Carlo simulations. Using this selection function, we have computed the Bright SHARC XCLF and find it to be fully consistent with a non-evolving XCLF to $L_x \simeq 5 \times 10^{44} \text{ erg s}^{-1}$ and $z \simeq 0.7$.

At $L_x > 5 \times 10^{44} \text{ erg s}^{-1}$, we find evidence for a deficit of clusters compared to that expected from a non-evolving XCLF. We detect only one such cluster in the redshift range $0.3 \leq z \leq 0.7$ when we would expect 4.9 clusters based on the local XCLF of De Grandi et al. (1999). The statistical significance of this deficit is $\simeq 96\%$. To increase the statistical significance of this possible deficit, we have combined the Bright SHARC and the 160 deg² survey of Vikhlinin et al (1998a). This joint survey covers $\simeq 260 \text{ deg}^2$ and contains only one confirmed $0.3 \leq z \leq 0.7$, $L_x > 5 \times 10^{44} \text{ erg s}^{-1}$ cluster, while we would expect 7.6 such clusters based on the local XCLF (De Grandi et al. 1999). The statistical significance of the deficit in this joint survey increases to 99.5%. These results remain preliminary because of incompleteness in the optical follow-up and uncertainties in the local XCLF.

Subject headings: cosmology: observations — galaxies: clusters: general — galaxies: evolution — surveys — X-rays: general

¹Based on data obtained at the Kitt Peak National Observatory, European Southern Observatory, the Canada–France–Hawaii Telescope and Apache Point Observatory

1. Introduction

The observed evolution of the space density of clusters of galaxies provides a powerful constraint on the underlying cosmological model. Many authors have demonstrated – both analytically and numerically – that the expected abundance of clusters, as a function of cosmic epoch, is a sensitive test of the mean mass density of the universe (Ω_m) and the type of dark matter (Press & Schechter 1974; Lacey & Cole 1993, 1994; Oukbir & Blanchard 1992, 1997; Henry 1997; Eke et al. 1996, 1998; Viana & Liddle 1996, 1999; Bryan & Norman 1998; Reichart et al. 1999; Borgani et al. 1999).

Measurements of the evolution of the cluster abundance have made significant progress over the past decade. For example, in their seminal work, Gioia et al. (1990) and Henry et al. (1992) computed the luminosity function of X-ray clusters extracted from the *Einstein* Extended Medium Sensitivity Survey (EMSS) and concluded that the X-ray Cluster Luminosity Function (XCLF) evolved rapidly over the redshift range of $0.14 \leq z \leq 0.6$.

The launch of the ROSAT satellite heralded a new era of investigation into the XCLF. The ROSAT All-Sky Survey (RASS) has provided new determinations of the local XCLF and has demonstrated that there is little observed evolution in the XCLF out to $z \sim 0.3$ (Ebeling et al. 1997; De Grandi et al. 1999) in agreement with the earlier work of Kowalski et al. (1984). In addition, the ROSAT satellite has supported several investigations of the distant X-ray cluster population (RIXOS, Castander et al. 1995; SHARC, Burke et al. 1997, Romer et al. 1999; RDCS, Rosati et al. 1998; WARPS, Jones et al. 1998; Vikhlinin et al. 1998a; NEP, Henry et al. 1998). Initially, such investigations reported a deficit of high redshift, low luminosity clusters consistent with the original EMSS result (Castander et al. 1995). However, over the last few years, there has been a growing consensus for a non-evolving XCLF. First, Nichol et al. (1997) re-examined the EMSS cluster sample and determined that the statistical evidence for evolution of the EMSS XCLF had decreased in light of new ROSAT data. Second, several authors have now conclusively shown that the XCLF does not evolve out to $z \sim 0.7$ for cluster luminosities of $L_x < 3 \times 10^{44} \text{erg s}^{-1}$ (Collins et al. 1997; Burke et al. 1997; Rosati et al. 1998; Jones et al. 1998).

Above $L_x = 3 \times 10^{44} \text{erg s}^{-1}$, recent work has indicated that the XCLF may evolve rapidly in agreement with the original claim of Gioia et al. (1990). Reichart et al. (1999) highlighted a deficit of luminous ($L_x > 5 \times 10^{44} \text{erg s}^{-1}$) EMSS clusters at $z > 0.4$ *i.e.* the EMSS survey has both the sensitivity and area to find such clusters but does not detect them. Moreover, Vikhlinin et al. (1998b) has recently reported evidence for a deficit of luminous clusters at $z > 0.3$ based on the 160 deg^2 ROSAT survey (Vikhlinin et al. 1998a).

In this paper, we report on the first determination of the bright end of the XCLF that is independent of the EMSS. In sections 2 & 3, we outline the Bright SHARC sample of clusters used herein and its selection function. In sections 4 & 5, we present the derivation of the XCLF and discuss its implications. Throughout this paper, we use $H_0 = 50 \text{ km s}^{-1} \text{ Mpc}$ and $q_0 = \frac{1}{2}$ to be consistent with other work in this field. All quoted luminosities are in the hard ROSAT passband [$0.5 \rightarrow 2.0 \text{ keV}$] and are aperture and k-corrected (see Romer et al. 1999 for details).

2. The Bright SHARC Sample

The details of the construction of the Bright SHARC survey are presented in Romer et al. (1999). The Bright SHARC was constructed from 460 deep ($T_{\text{exp}} > 10 \text{ ksecs}$), high galactic latitude ($|b| > 20^\circ$), ROSAT

PSPC pointings which cover a unique area of 178.6 deg^2 . Using a wavelet-based detection algorithm, 10277 X-ray sources were detected in these pointings of which 374 were measured to be significantly extended ($> 3\sigma$; see Nichol et al. 1997) relative to the ROSAT PSPC point-spread function. The Bright SHARC represents the brightest 94 of these 374 extended cluster candidates above a ROSAT count rate of $0.0116 \text{ cnts s}^{-1}$. This corresponds to a flux limit of $\simeq 1.4 \times 10^{-13} \text{ erg s}^{-1} \text{ cm}^2$ [0.5–2.0 keV] for the average neutral hydrogen column density of the Bright SHARC and a cluster temperature of 6 keV.

Over the past two years, we have optically identified the most likely X-ray emitter for 91 of these 94 Bright SHARC cluster candidates and have discovered 37 clusters, 3 groups of galaxies and 9 nearby galaxies (the remainder are blends of X-ray sources *e.g.* AGNs & stars; see Romer et al. 1999). We find 12 clusters in the range $0.3 \leq z \leq 0.83$ (median redshift of $z = 0.42$) and have independently detected cluster RXJ0152-7363 ($z = 0.83$ based on 3 galaxy redshifts obtained at the CFHT) which is one of the most luminous, high redshift X-ray clusters ever detected (see Romer et al. 1999). This cluster has also been detected by the WARPS and RDCS surveys (see Ebeling et al. 1999; Rosati, private communication).

3. Selection Function

An important part of any survey is a solid understanding of the selection function *i.e.* the efficiency of finding objects as a function of both cosmological and operational parameters. In the case of the EMSS cluster sample, the selection function is somewhat straightforward since the EMSS optically identified all sources regardless of their observed X-ray extent. This is not the case for the Bright SHARC and therefore, the most direct way of modelling the selection function is through Monte Carlo simulations. The details of such simulations are given in Adami et al. (1999) but we present here some initial results.

The Bright SHARC selection function is obtained by adding artificial clusters to PSPC pointings and determining if these clusters would have satisfied the Bright SHARC selection criteria. In this way, we can accurately model the effects of the signal-to-noise cut, the extent criteria and blending of sources. To date, we have only simulated artificial clusters at four different intrinsic luminosities, $L_x = 1, 2, 5 \text{ \& } 10$ ($\times 10^{44} \text{ erg s}^{-1}$ *i.e.* equally spaced in log-space), and covering a redshift range of $0.3 \leq z \leq 1$ in steps of $\delta_z = 0.05$. These artificial clusters were constructed assuming an isothermal King profile with $r_c = 250 \text{ kpc}$ and $\beta = \frac{2}{3}$. The effects of changing the cluster profile and its parameters are explored in detail in Adami et al. (1999).

For each combination of L_x and z , we added, one at a time, 10 artificial clusters to each of 10 PSPC pointings (randomly chosen from all pointings available in the SHARC survey). The positions of the clusters in these pointings were chosen at random but recorded for later use when computing the area surveyed (see below). For each artificial cluster, we computed the expected number of ROSAT PSPC counts for its given luminosity, redshift and the exposure time of the pointing. We then took a Poisson deviate about the expected number of counts and distributed the counts at random assuming a redshifted King profile. The pointing was then processed in the same fashion as the real data (see Romer et al. 1999; Nichol et al. 1997) thus allowing us to determine if the cluster would have been selected for the Bright SHARC.

In Figure 1, we show the results of these initial simulations. The effective area of the Bright SHARC was computed by splitting the PSPC field-of-view into 4 annulii ($2.5 \rightarrow 6.25$, $6.25 \rightarrow 11.25$, $11.25 \rightarrow 16.25$ and $16.25 \rightarrow 22.5$; the central part of the PSPC was excluded) and multiplying the area in each of these annulii by the measured success rate of detecting clusters in these same annulii (as a function of the input cluster redshift and luminosity). We then summed the effective area in these annulii over all pointings used –




Fig. 1.— The effective area of the Bright SHARC sample of clusters as a function of cluster luminosity and redshift. These curves were constructed from the initial results of our Monte Carlo simulations (see Adami et al. 1999). The four curves represent the four different input luminosities (in units of $10^{44} \text{ erg s}^{-1}$). The simulations were performed over a redshift range of $0.3 \leq z \leq 1$ in steps of $\delta_z = 0.05$.

again as a function of luminosity and redshift – to provide the total effective area sampled by the simulations. Finally, we re-scaled the results to obtain the expected area for all 460 PSPC pointings used in the Bright SHARC.

4. Luminosity Function

The luminosity function of the Bright SHARC was determined using the $1/V_a$ methodology outlined in Avni & Bahcall (1980), Henry et al. (1992) and Nichol et al. (1997), where V_a is the available sample volume for any given cluster in the survey. Using the selection function presented in Figure 1, V_a can be computed for a cluster of luminosity L_x using

$$V_a = \int_{z_{low}}^{z_{high}} \Omega(L_x, z) V(z) dz, \quad (1)$$

where z_{low} and z_{high} are the lower and upper bounds of the redshift shell of interest, $V(z)$ is the volume per unit solid angle for that redshift shell and $\Omega(L_x, z)$ is the effective area of the Bright SHARC from Figure 1. In practice, the integral in Eqn. 1 is replaced by a sum over the discrete values of $\Omega(L_x, z)$ obtained from




Fig. 2.— The XCLF for the Bright SHARC survey (solid stars) and the Southern SHARC (solid circles; taken from Table 1 of Burke et al. 1997). The y-axis was computed using Equation 2 with ΔL values quoted in Table 1 and has units of $\text{Mpc}^{-3}(10^{44} \text{ erg s}^{-1})^{-1}$. These data points span a redshift range of $0.3 \leq z \leq 0.7$. The hashed region represents the local XCLF as measured by De Grandi et al. (1999) and Ebeling et al. (1997); we have included the $\pm 1\sigma$ errors as given by these authors. The unfilled stars are taken from Nichol et al. (1997) and are the EMSS luminosity function at $0.3 \leq z \leq 0.6$ including new data from the ROSAT pointing archive. The solid triangle is the XCLF expected for RXJ0152-7363 and its value is presented in Table 1. All the error bars are computed using the Poisson errors of Gehrels (1986)

the simulations. Linear interpolation was used where necessary to obtain finer resolution in both luminosity and redshift space.

The luminosity function was derived by summing the V_a values for all clusters in the Bright SHARC as a function of luminosity *i.e.*

$$n(L) = \frac{1}{\Delta L} \sum_{i=1}^N \frac{1}{V_a^i}, \quad (2)$$

where ΔL is the width of the luminosity bins and N is the number of clusters in that luminosity bin (Table 1). For the results presented here, we have restricted ourselves to the luminosity range $10^{44} \leq L_x \leq 10^{45} \text{ erg s}^{-1}$.

In Figure 2, we present the Bright SHARC XCLF and compare it to the Southern SHARC XCLF (Burke et al. 1997) and measurements of the local XCLF (Ebeling et al. 1997; De Grandi et al. 1999). We provide,

Table 1: The measured space density of the Bright SHARC sample of clusters

redshift shell	$\log_{10} L_x$ [bin center]	$\log_{10} n(L)$ [$\text{Mpc}^{-3}(10^{44}\text{erg s}^{-1})^{-1}$]	$\log_{10} \Delta L$ [10^{44}erg s^{-1}]	No. clusters
$0.3 \leq z \leq 0.7$	44.23	-6.74	1.4	3
$0.3 \leq z \leq 0.7$	44.49	-7.39	1.4	6
$0.3 \leq z \leq 0.7$	44.65	-8.09	1.4	2
$0.3 \leq z \leq 1.0$	44.93	-8.99	3	1

in Table 1, the data points displayed in Figure 2 together with the redshift and luminosity ranges studied. We also provide the number of clusters in each bin. We have not performed a parametric fit to the data because of the limited dynamic range in luminosity available from our present simulations.

5. Discussion

Figure 2 demonstrates that the high redshift XCLF does not evolve below $L_x = 5 \times 10^{44}\text{erg s}^{-1}$, or below L^* in the XCLF ($L^* = 5.7^{+1.29}_{-0.93} \times 10^{44}\text{erg s}^{-1}$ from Ebeling et al. 1997). Using a Kolmogorov–Smirnov (KS) test similar to that discussed by De Grandi et al. (1999), we find that the Bright SHARC unbinned data is fully consistent with the low redshift XCLF over the luminosity range $1 \leq L_x \leq 5 \times 10^{44}\text{erg s}^{-1}$ (we find a KS probability of 0.32). This result is consistent with previous work (Nichol et al. 1997; Burke et al. 1997; Rosati et al. 1998; Jones et al. 1998) but pushes the evidence for a non-evolving XCLF to the highest luminosities presently reached by ROSAT data. This is illustrated by the fact that we find 6 distant clusters in the luminosity range $3 \leq L_x \leq 5 \times 10^{44}\text{erg s}^{-1}$ (Table 1) which is more than any other ROSAT archival survey. The issue therefore becomes the degree of observed evolution in the XCLF above L^* .

We have investigated evolution in the XCLF above L^* in two separate ways. First, we have one very luminous high redshift cluster in the Bright SHARC – RXJ0152.7-1357 ($z = 0.83$; $L_x = 8.5 \times 10^{44}\text{erg s}^{-1}$) – which probes the XCLF above L^* . The implied position of this cluster in the XCLF is given in Figure 2 and Table 1 (we used a redshift range of $0.3 \leq z \leq 1$ and luminosity range of $7 \times 10^{44} \leq L_x \leq 10^{45}\text{erg s}^{-1}$ when computing the volume sampled by this cluster). As can be seen in Figure 2, the implied space density of RXJ0152-1357 agrees with the local XCLF and may be evidence for a non-evolving XCLF above L^* . However, we must remain cautious since RXJ0152.7-1357 has a complex X-ray morphology indicative of an on-going merger which may have enhanced its luminosity (see Ebeling et al. 1999). Such disturbed or non-spherical morphologies (both in the X-rays and optical) appear to be common at these high redshifts – *e.g.* MS1054.4-0321 at $z = 0.823$ (Donahue et al. 1999) and RXJ1716.6+6708 at $z = 0.813$ (Henry et al. 1998; Gioia et al. 1999) – and may indicate that we are witnessing the epoch of massive cluster formation.

The second path of investigation is to repeat the analysis of Collins et al. (1997) and Vikhlinin et al. (1998b) and compute the number of expected Bright SHARC clusters at these bright luminosities assuming a non-evolving XCLF. In the luminosity range $5 \times 10^{44} \leq L_x \leq 10^{45}\text{erg s}^{-1}$ and redshift range $0.3 \leq z \leq 0.7$, we would predict 4.9 clusters based on the De Grandi et al. (1999) XCLF (or 3.5 clusters using the Ebeling et al 1997 XCLF). At present, the Bright SHARC contains only one confirmed cluster in this range, RXJ1120.1+4318 at $z = 0.60$ and $L_x = 5.03 \times 10^{44}\text{erg s}^{-1}$. The Poisson statistical significance (Gehrels 1986) of this observed deficit is $\simeq 96\%$ (or $\simeq 90\%$ for Ebeling et al. 1997).

One way to increase the statistical significance of any possible deficit of high redshift X-ray luminous clusters is to combine the Bright SHARC and 160deg² survey of Vikhlinin et al. (1998a) as both surveys should have similar selection functions. We have determined that 201 ROSAT PSPC pointings are in common between the two surveys (see Romer et al. 1999), or $\simeq 44\%$ of the area of the Bright SHARC. Vikhlinin et al. (1998b) have noted that their survey probably contains only 2 candidate clusters above $L_x = 3 \times 10^{44} \text{ erg s}^{-1}$ and $z > 0.3$ (based on photometric redshift estimates), while they would have expected 9 clusters above this redshift and luminosity limit. In fact, Bright SHARC spectroscopy of one of these two candidate clusters – RXJ1641+8232 – shows that it is at $z = 0.195$ (based on 3 galaxy redshifts). The other candidate cluster – RXJ1641+4001 – is not in the Bright SHARC.

Therefore, the Bright SHARC plus the 160deg² survey cover a total area of $\sim 260 \text{ deg}^2$ and scaling the aforementioned numbers appropriately, we would expect 7.6 (using the De Grandi et al. 1999 XCLF) or 5.5 (using the Ebeling et al. 1997 XCLF) clusters in the luminosity range $5 \times 10^{44} \leq L_x \leq 10^{45} \text{ erg s}^{-1}$ and redshift range $0.3 \leq z \leq 0.7$. By comparison, this joint survey contains only 1 confirmed² X-ray luminous, high redshift cluster. From Gehrels (1986), the statistical significance of this deficit is now 99.5% or 97.5% respectively. We note however that there are still candidates in both surveys that require further optical follow-up *i.e.* one of the three unidentified Bright SHARC candidates mentioned in section 2 could be a high redshift cluster.

The above analyses suffer from small number statistics and incomplete optical follow-up. Moreover, the local XCLFs are also affected by small number statistics at these high X-ray luminosities. Therefore, the issue of evolution above L^* in the XCLF remains unclear. However, we note that four independent surveys of distant clusters of galaxies have now seen a potential deficit of X-ray luminous high redshift clusters (SHARC, WARPS, EMSS, 160 deg² survey). It is worth stressing that some level of XCLF evolution is expected above L^* at high redshift and the degree of such evolution is a strong indicator of Ω_m (see Blanchard & Oukdir 1992, 1997; Reichart et al. 1999).

The way to resolve these problems is to construct larger samples of clusters over bigger areas of the sky. This is certainly possible for the local XCLF *e.g.* using the on-going REFLEX survey of clusters being constructed from the RASS (see Böhringer et al. 1998). However, for the distant XCLF, it is unlikely that a significant amount of further area ($>> 260 \text{ deg}^2$) can be added to the present surveys using the existing ROSAT pointing archive; one may have to wait for the XMM satellites (see Romer 1998). The next major improvements to the results presented in this paper will be to obtain a more detailed view of the Bright SHARC selection function as well as to complete the optical follow-up of the remaining candidates. Another improvement would be the possible combination of all the ROSAT distant surveys (SHARC, 160 deg², RDCS, NEP & WARPS) thus maximising the amount of volume sampled at high redshift.

6. Acknowledgements

The authors would like to thank Alain Blanchard, Jim Bartlett, Francisco Castander, Harald Ebeling, Pat Henry, Andrew Liddle, Piero Rosati & Alex Vikhlinin for helpful discussions over the course of this work. This research was supported through NASA ADP grant NAG5-2432 (at NWU) and NASA LTSA grant NAG5-6548 (at CMU). AM thanks the Carnegie Mellon Undergraduate Research Initiative for financial

² On-going spectroscopic redshift measurements of further candidate clusters in the 160deg² survey have yet to reveal a single $L_x > 5 \times 10^{44} \text{ erg s}^{-1}$ cluster (Vikhlinin, private communication)

assistance. We thank Alain Mazure and the IGRAP–LAS (Marseille, France) for their support.

REFERENCES

- Adami, C., Ulmer, M. P., Pildis, R. A., Romer, A. K., Holden, B. P., & Nichol, R. C., 1999, *ApJ*, in preparation
- Avni, Y., Bahcall, J. N., 1980, *ApJ*, 235, 694
- Böhringer, H., Guzzo, L., Collins, C.A., Neumann, D.M., Schindler, S., Schuecker, P., Cruddace, R., Chincarini, G., De Grandi, S., Edge, A.C., Macgillivray, H.T., Shaver, P., Vettolani, G., Voges, W. 1998, in *Wide Field Surveys in Cosmology*, Editions Frontieres. ISBN 2-8 6332-241-9, 261.
- Borgani, S., Rosati, P., Tozzi, P., Norman, C. 1999, in press (astro-ph/9901017).
- Bryan, G. L., Norman, M. L., 1998, *ApJ*, 495, 80
- Burke, D.J., Collins, C.A., Romer, A.K., Holden, B.P., Nichol, R.C. 1997, *ApJ*, 488, L83.
- Castander, F.J., Bower, R.G., Ellis, R.S., Aragon-Salamanca, A., Mason, K.O., Hasinger, G., McMahon, R.G., Carrera, F.J., Mittaz, J.P.D., Perez-Fournon, I., Lehto, H.J. 1995, *Nature*, 281, 59.
- Collins, C.A., Burke, D.J., Romer, A.K., Sharples, R.M., Nichol, R.C. 1997, *ApJ*, 479, L117.
- De Grandi, S., Böhringer, H., Guzzo, L., Molendi, S., Chincarini, G., Collins, C.A., Cruddace, R., Neumann, D.M., Schindler, S., Schuecker, P., & G., Voges, W., 1999, *ApJ*, 514, in press (astro-ph/9902067)
- Donahue, M., Voit, G.M., Gioia, I., Luppino, G., Hughes, J.P., Stocke, J.T. 1998, *ApJ*, 502, 550.
- Ebeling, H., Edge, A. C., Fabian, A. C., Allen, S., Crawford, C. S., Böhringer, H., 1997, *ApJ*, 479, L101
- Ebeling, H., Edge, A.C., Böhringer, H., Allen, S.W., Crawford, C.S., Fabian, A.C., Voges, W., Huchra, J.P. 1998, *MNRAS*, 301, 881.
- Ebeling, H., Jones, L.R. , Perlman, E., Scharf, C., Horner, D., Wegner, G., Malkan, M., Fairley, B., Mullis, C. R., 1999, *ApJ*, submitted (astro-ph/9905321) Eke, V. R., Cole, S., Frenk, C., 1996, *MNRAS*, 282, 263
- Eke, V. R., Cole, S., Frenk, C., & Henry, J. P., 1998, *MNRAS*, 298, 1145
- Gehrels, N. 1986, *ApJ*, 303, 336
- Gioia, I. M., Henry, J.P., Maccacaro, T., Morris, S.L., Stocke, J.T., Wolter, A. 1990, *ApJ*, 356, L35.
- Gioia, I. M., Henry, J.P., Mullis, C. R., Ebeling, H., Wolter, A. 1999, *ApJ*, accepted (astro-ph/9902277).
- Henry, J.P., Gioia, I.M., Maccacaro, T., Morris, S.L., Stocke, J.T., Wolter, A. 1992, *ApJ*, 386, 408.
- Henry, J.P. 1997, *ApJ*, 489, L1.
- Henry, J.P., Gioia, I.M., Mullis, C. R., Clowes, D. L., Luppino, G. A., Böhringer, H., Briel, U. G., Voges, W., Huchra, J. P., 1997, *AJ*, 114, 1293
- Jones, L.R., Scharf, C., Ebeling, H., Perlman, E., Wegner, G., Malkan, M., Horner, D. 1998, *ApJ*, 495, 100.
- Kowalski, M. P., Ulmer, M. P., Cruddace, R. G., Wood, K. S., 1984, *ApJ Supp.*, 56, 403.
- Lacey, C., Cole, S. 1993, *MNRAS*, 262, 627
- Lacey, C., Cole, S. 1994, *MNRAS*, 271, 676
- Nichol, R.C., Holden, B.P., Romer, A.K., Ulmer, M.P., Burke, D.J., Collins, C.A., *ApJ* 1997, 481, 644.
- Oukbir, J., Blanchard, A., 1992, *A&A*, 262, 21
- Oukbir, J., Blanchard, A., 1997, *A&A*, 317, 365
- Press, W. H., Schechter, P. 1974, *ApJ*, 187, 425
- Reichart, D.E., Nichol, R.C., Castander, F.J., Burke, D.J., Romer, A.K., Holden, B.P., Collins, C.A., Ulmer, M.P., *ApJ* 1999, accepted (astro-ph/9802153).
- Romer, A. K., 1998, in *Wide Field Surveys in Cosmology*, Editions Frontieres. ISBN 2-8 6332-241-9, 261. (astro-ph/9809198)
- Romer, A.K., Nichol, R.C., Holden, B.P., Ulmer, M.P., Pildis, R.A., Merrelli, A. J., Adami, C. A., Burke,

- D.J., Collins, C.A., Metevier, A., Kron, R. Commons, K., 1999, ApJS, submitted.
- Rosati, P., Della Ceca, R., Norman, C., Giacconi, R., 1998, ApJ, 492, 21.
- Viana, P.T.P., Liddle, A.R. 1996, MNRAS, 281, 323
- Viana, P.T.P., Liddle, A.R. 1999, in press (astro-ph/9803244)
- Vikhlinin, A., McNamara, B.R., Forman, W., Jones, C., Quintana, H., Hornstrup, A. 1998a, ApJ 502, 558.
- Vikhlinin, A., McNamara, B.R., Forman, W., Jones, C., Quintana, H., Hornstrup, A. 1998b, ApJ 498, L21.